



General Instructions

General: You can expect to need 1 week(s) off from work.

Pre-operative:

1. Three weeks before surgery, make sure you have discontinued all medications that are associated with increased bleeding. These include aspirin, ibuprofen, aleve, Plavix, Coumadin/warfarin, vitamin E, fish oil, xarelto, eliquis. If you are on any of these medications you must let Dr. Wilson and his staff know. You may take Tylenol and extra strength Tylenol only.
2. Patients must be non-smokers. If you are a smoker you must have stopped smoking for six weeks before any surgical procedures will be performed. Nicotine products may be used to aid in smoking cessation, but must also be stopped prior to surgery. You may be asked to complete a blood test to check the circulating levels of nicotine in your system prior to surgery.
3. We recommend you purchase arnica tablets and begin them one week prior to your procedure and continue for the week following the procedure. These are available at our online dispensary or at your local health food store.
4. You must have someone drive you to and from your surgery. You must have an adult to help you for the first 24 hours. If you do not have someone to help you, the office can help you coordinate a nurse or assistant.
5. Dr. Wilson will prescribe you your pain medication the day of the procedure. Rarely will you need narcotic pain medications for this procedure.
6. **For hospital procedures:** Do not eat or drink anything after midnight the night before surgery. If you have a medication you must take in the morning, take with a small sip of water. Shower the morning of surgery making sure there are no traces of makeup left on your skin. Do not apply moisturizer or anything to your face the morning of surgery and do not wear contact lenses. Please wear comfortable clothing that either zips or buttons in the front.
7. **For In Office Procedures:** If prescriptions are required they will be sent to your pharmacy prior to procedure. Take medications as directed. If taking narcotic pain medications you will require a ride to and from your procedure. Local anesthetic may be used.
8. If you have hair long enough, please place your hair in a tight ponytail the day of your surgery.



General Instructions

Post Operative: First 48 Hours

1. Notify Dr. Wilson if you have any increased bruising or asymmetrical swelling. This may be a sign of bleeding.
2. Do not get the area wet for the first 48 hours. After this time you may shower, but avoid direct water pressure to the area. Do not scrub or soak the area.
3. Keep any exposed sutures/suture-lines covered with bacitracin ointment. Gently wipe and re-apply bacitracin twice daily.
4. If you have steri strips in place you do not need to apply bacitracin ointment. The steri strips can be gently removed after one week.
5. Take pain medications as prescribed. Only take the medications as needed. Do not take ibuprofen, aspirin or Aleve.
6. Do not use ice unless specifically instructed by Dr. Wilson or his staff.
7. Resume a regular diet. If the first night you do not feel up to eating, ensure you take in plenty of fluids. Do not take medications on an empty stomach.
8. Keep Bacitracin ointment on the surgical area. This should be applied twice daily. You may gently wipe off prior to reapplying.

Post Operative: 2 days to 2 weeks

1. Your stitches will dissolve they will not need to be removed. Place bacitracin ointment on the site until the stitches all dissolve. You can shower 48 hours after your surgery. Do not scrub incisions. Let soapy water gently glide over sites.
2. Resume a regular complete diet.
3. Refrain from exercise for two weeks at which point you can start light activities. No heavy exercise for six weeks.
4. You can resume your Aspirin and other medications associated with bleeding risk one week after your procedure.
5. Do not hesitate to call the office with any questions. In case of an emergency please contact the office at 603-294-4526.